



Bergvesenet rapport nr <b>BV BV 2818</b>	Intern Journal nr	Gammelt internt rapp. nr.	Rapport lokalisering Trondheim	Gradering <b>Apen</b>
Kommer fra ..arkiv	Ekstern rapport nr Aspro 1506	Oversendt fra Bidjovagge Gruber a. s.	Fortrolig pga	Fortrolig fra dato:

**Tittel**

Detailed geological suvey on electromagnetic anomalies.  
 Kautokeino, West-Finmark

Forfatter Berge, Kari	Dato    År 27/4 1984	Bedrift Prospektering A/S
--------------------------	-------------------------	------------------------------

Kommune Kautokeino	Fylke Finmark	Bergdistrikt	1: 50 000 kartblad 18332 19324	1: 250 000 kartblad
-----------------------	------------------	--------------	--------------------------------------	---------------------

Fagområde Geologi Geofysikk	Dokument type Rapport	Forekomster
Råstoffgruppe Malm/metall	Råstofftype Fe Grafitt	

**Sammendrag / innholdsfortegnelse**

The survey was made to examine electromagnetic anomalies from Digheim's airborne survey of the Superior Oil Joint Venture Area.

In this report 13 areas are described. The location of the area are shhown on figs. no. 1, 2 and 3.

Detailed mapping has been done within grids put up for the ground geophysics.

The areas are for the major part covered by moraine. In addition to bedrock mapping, boulders were also mapped. Possible causes for the electromagnetic anomalies were especially looked for, such as zones of iron sulphides, iron oxides and graphite.

The results are presented on topographical maps in scale 1:5000, Figs. no.4 to 15. The area numbers correspond to the numbers which the areas are given in the ground geophysics survey.

Together with the geology, the maps also show electromagnetic profiles from the ground geophysics survey.

To each map there is a description og geology, boulders and mineralizations.

Each area is given a recomandation for further follow -up work. This follow-up work was done partly in 1983 and will be continued in 1984.



INTERN RAPPORT.

GAMLE RINGERIKS VEI 14 POSTB. 83 - 1321 STABEKK  
MELEID AV AKTIESELSKABET BYDVARANGER

Telex 72 987 aspro n

DATO: 27/4-84

RAPPORT NR: 1506

KARTBLAD 1833 II  
1932 IV

Antall sider  
— " — bilag

SAKSBEARBEIDER Kari Berge

RAPPORT VEDRØRENDE:

Detailed geological survey on electromagnetic anomalies.  
Kautokeino, West-Finnmark

FORDELING  
OSLO:

Vertical grid of 10 empty boxes for distribution in Oslo.

RESYMÉ:

The survey was made to examine electromagnetic anomalies from Dighem's airborne survey of the Superior Oil Joint Venture Area.

In this report 13 areas are described. The location of the areas are shown on figs. no. 1, 2 and 3.

Detailed mapping has been done within grids put up for the ground geophysics. The areas are for the major part covered by moraine. In addition to bedrock mapping, boulders were also mapped. Possible causes for the electromagnetic anomalies were especially looked for, such as zones of iron sulphides, iron oxides and graphite.

KIRKENES:

Vertical grid of 8 empty boxes for distribution in Kirkenes.

ANDRE:

Vertical grid of 8 empty boxes for distribution elsewhere.

The results are presented on topographical maps in scale 1:5000, Figs. no. 4 to 15. The area numbers correspond to the numbers which the areas are given in the ground geophysics survey. Together with the geology, the maps also show electromagnetic profiles from the ground geophysics survey. To each map there is a description of geology, boulders and mineralizations.

Each area is given a recommendation for further follow-up work. This follow-up work was done partly in 1983 and will be continued in 1984.

KOMMENTAR:




Fig. 1: Location map




Fig. 2: Location map




Fig. 3: Location map

AREA 30

AREA 30 Fig. 4

The localization of the area is uncertain. No sign of the grid except for one unmarked pole, was found.

Boulders

Carbonate breccia and carbonate rock with quartz veins.

Coarse grained amphibolite or diabase, and amphibolite.






Fine grained tuff or tuffite with veins filled with quartz and carbonate.

Recommendations

The geological observations give little information. The anomaly is suggested further examined by deep till sampling.

LEGEND

Geology

-  Granite / P=pegmatite
-  Mica schist
-  Coarse, massive amphibolite or diabase
-  Tuff, tuffite
-  Medium to finegrained, often schistose and foliated, amphibolite
-  Carbonate rock, albite carbonate rock
-  Carbonate schist
-  Carbonate breccia
-  Sandstone, quartzite
-  Gneiss
-  Rust
-  Strike and dip (90 degrees division)
-  Rock boundary, certain and uncertain
-  Outcrop
-  Boulder
-  Group of boulders



hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopryite


} As a major constituent (> 1/3)  
of the rock

Electromagnetics

-  Real comp Horizontal loop 1777 Hz
-  Imag comp Horizontal loop 1777 Hz

Topography

-  Stream or river, lake
-  Bog
-  Elevation contour
-  Elevation 505
-  Path
-  Cart track
-  UTM grid with coordinates



492  
x

AREA 30 Kautokeino Geological mapping EM survey	<b>M</b> 1:5000
Målt: KB 63	Tegn: KB 3/64
Trace: HB 4/64	Fig. 4
<b>PROSPEKTERING A/S</b>	

AREA 31

AREA 31 Fig. 5


A part of the area is investigated. No outcrops were discovered.

Except for some boulders of gneiss and amphibolite, there are mainly boulders of albite carbonate rock. These are very fine grained, light coloured and have veins with hematite mineralization.

Recommendations






As there is little geological informations on this area, the anomaly is suggested surveyed in a minor deep till sampling programme.

500 W 400 W 300 W 200 W 100 W 0 100 E 200 E



LEGEND

Geology

-  Granite / P=pegmatite
-  Mica scist
-  Coarse, massive amphibolite or diabase
-  Tuff, tuffite
-  Medium to fine grained, often schistose and foliated, amphibolite
-  Carbonate rock, albite carbonate rock
-  Carbonate schist
-  Carbonate breccia
-  Sandstone, quartzite
-  Gneiss
-  Rust
-  Strike and dip (90 degrees division)
-  Rock boundary, certain and uncertain
-  Outcrop
-  Boulder
-  Group of boulders




hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopryrite

} As a major constituent (> 1/3)  
 of the rock

Electromagnetics

-  Real comp. Horizontal loop 1777 Hz
-  Imag comp. Horizontal loop 1777 Hz

Topography

-  Stream or river, lake
-  Bog
-  Elevation contour
-  Elevation
-  Path
-  Cart track
-  UTM grid with coordinates

AREA 31 Kautokeino Geological mapping EM survey	M
	1:5000
	Mått: KB 63
	Tegn: KB 3/84
	Trace: HB 4/84
PROSPEKTERING A/S	Fig.5

AREA 32

AREA 32 Fig. 6


Inside the grid area there are no exposures of bedrock. About 200 m east of the grid there is an outcrop of tuff or tuffite. It has veins filled with carbonate minerals and quartz. The rock is probably altered.

Boulders

Boulders of carbonate schist are dominating the area. There are also boulders of coarse grained amphibolite and pegmatitic granite.

Recommendations

There is little geological information on this area. The anomaly is recommended surveyed by deep till sampling.



**LEGEND**

Geology

- Granite / P=pegmatite
- Mica schist
- Coarse, massive amphibolite or diabase
- Tuff, tuffite
- Medium to fine grained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust
- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 ml - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3) of the rock

Electromagnetics

- Real comp. Horizontal loop 1777 Hz
- Imag. comp. Horizontal loop 1777 Hz

Topography

- Stream or river, lake
- Bog
- Elevation contour
- Elevation
- Path
- Cart track
- UTM grid with coordinates

Area 32 Kautokeino Geological mapping EM survey	M 1:5000
	Målt: KB 83
	Tegn: KB 3/84
	Trace: HB 4/84
PROSPEKTERING A/S	Fig. 6


AREA 33

AREA 33 Fig. 7

The area lacks exposures of bedrock. It is covered by bog and moraine. Boulders that are seen consist of banded gneiss, carbonate breccia, carbonate schist, tuff or tuffite and quartzite.

Recommendations

No further work is recommended in this area.



**LEGEND**

**Geology**

- Granite / P=pegmatite
- Mica schist
- Coarse, massive amphibolite or diabase
- Tuff, tuffite
- Medium to finegrained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust
- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3) of the rock

**Electromagnetics**

- Real comp. Horizontal loop 1777 Hz
- Imag comp. Horizontal loop 1777 Hz

**Topography**

- Stream or river, lake
- Bog
- Elevation contour
- Elevation
- Path
- Cart track
- UTM grid with coordinates

AREA 33 Kautokeino Geological mapping EM survey	M
	1:5000
	Målt: KB 83
	Tegn: KB 3/84
	Trace: HB 4/84
PROSPEKTERING A/S	Fig. 7

AREA 34

AREA 34 Fig. 8

The surveyed area is dominated by bog and moraine.

Outcrops

Along the 100W profile there are outcrops of albite carbonate rocks and carbonate breccia.

North of the area there are outcrops as far north as the lake Bædnatjav'ri. On both sides of a small stream that leads southwards from the lake towards the measured area, there are outcrops of coarse grained quartzite. The quartzite is exposed in topographic depressions.

To the east of the quartzite, and seemingly overlying this, there are albite carbonate rocks. These consist of zones or layers of differing varieties of the rock: One light coloured to reddish and very fine grained rock with hematite in grains with sizes up to 1 cm across, and one grey, fine grained rock containing some mica. The zones occur irregularly and the lighter type may be an alteration product of the grey one.

One outcrop is of an amphibolitic rock type.


The rocks are mostly without orientation. Two outcrops show northwestern strike and have a steep and moderate dip to the southwest.

Boulders

The moraine in the eastern part of the surveyed area contains boulders of granite, gneiss, amphibolite, albite carbonate rocks and quartzite.

Recommendations

Due to the geological observations the area is recommended investigated by diamond drilling.



**LEGEND**

**Geology**

- Granite / P=pegmatite
- Mica scist
- Coarse, massive, amphibolite or diabase
- Tuff, tuffite
- Medium to finegrained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust
- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3) of the rock

**Electromagnetics**

- Real comp. Horizontal loop 1777 Hz
- Imag comp. Horizontal loop 1777 Hz

**Topography**

- Stream or river, lake
- Bog
- Elevation contour
- Elevation 605
- Path
- Cart track
- UTM grid with coordinates 67 90

AREA 34 Kautokeino Geological mapping EM survey	M
	1:5000
	Målt: KB 63
	Tegn: KB 3/84
	Trace: HB 4/84
PROSPEKTERING A/S	Fig. 8

AREA 35

AREA 35 Fig. 9

The area has a thick cover of moraine.

Outcrops are found only by the riverside. There are outcrops of diabase, carbonate breccia and tuff.


Boulders

Where the moraine cover is thick, there are boulders mostly of gneiss and quartzite. Along the river bank there are boulders of granite, amphibolite and albite carbonate rocks as well.

Recommendations

Geological observations are lacking in most of the area.

To examine the anomaly further, deep till sampling is recommended.



LEGEND

Geology

- Granite / P=pegmatite
- Mica schist
- Coarse, massive amphibolite or diabase
- Tuff, tuffite
- Medium to fine grained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust

- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3) of the rock

Electromagnetics

- Real comp. Horizontal loop 1777 Hz
- Imag. comp. Horizontal loop 1777 Hz

Topography

- Stream or river, lake
- Bog
- Elevation contour
- Elevation 605
- Path
- Cart track
- UTM grid with coordinates

AREA 35 Kautokeino Geological mapping EM survey	M
	1:5000
	Målt: KB 83
	Tegn: KB 3/83
	Trace: HB 4/84
PROSPEKTERING A/S	Fig. 9

AREA 36

AREA 36 Fig. 10

There are no outcrops inside the grid area on the southern side of the river. The land rises steeply on the southern bank of the river to a thick cover of moraine.

Further west on the southern river bank, there are outcrops of coarse grained amphibolite and carbonate rock.


Boulders

Boulders in the moraine consist of gneiss for the major part.

Close to the river bank, the boulders are amphibolite and carbonate rock.

Recommendations

Although there is little information on the geology of this area, the anomaly is interesting and diamond drilling is recommended.



**LEGEND**

**Geology**

- Granite / P=pegmatite
- Mica schist
- Coarse, massive amphibolite or diabase
- Tuff, tuffite
- Medium to finegrained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust
- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 ml - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3) of the rock.

**Electromagnetics**

- Real comp. Horizontal loop 1777 Hz
- Imag. comp. Horizontal loop 1777 Hz

**Topography**

- Stream or river, lake
- Bog
- Elevation contour
- Elevation
- Path
- Cart track
- UTM grid with coordinates

AREA 36 Kautokeino Geological mapping EM survey	M 1:5000
	Målt: KB 83 Tegn: KB 3/84 Trace: HB 4/84
	PROSPEKTERING A/S Fig. 10

AREA 44

AREA 44 Fig. 11


The area is well exposed. Only one outcrop is examined in this investigation.

The rock is a quartz-mica schist. It is grey, foliated and fine grained. The foliation is cut by a vein of pegmatitic granite. The strike of the rock is to the southeast, and it is dipping gently to the southwest.

Recommendations

The geological information is small. On the strength of the anomaly, deep till sampling is recommended.

Area 44



LEGEND

Geology

- Granite / P=pegmatite
- Mica schist
- Coarse, massive amphibolite or diabase
- Tuff, tuffite
- Medium to finegrained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust
- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3)  
 of the rock

Electromagnetics

- Real comp. Horizontal loop 1777 Hz
- Imag comp. Horizontal loop 1777 Hz

Topography

- Stream or river, lake
- Bog
- Elevation contour
- Elevation
- Path
- Cart track
- UTM grid with coordinates

AREA 44 Lavvoaivi Geological mapping EM survey	M
	1:5000
	Målt: KB 63
	Tegn: KB 3/84
Trace: HB 4/84	Fig. 11
PROSPEKTERING A/S	

AREA 4546

## AREA 45 Fig. 12

The bedrock in the area is well exposed. The rocks are foliated amphibole schists with lenses and layers of quartzite. There are also outcrops of granite and gneiss. To the far north in the area there is amphibole biotite schist that is mapped as mica schist. Coarse and massive amphibole schist is also separated from the amphibole schist on the map.

### Amphibole schist

The rock is foliated with bands of amphibole, feldspar and various amounts of quartz. The bands are about 1 mm in width. Zones that are rich in quartz frequently show distorted foliation and lack measurable orientation. The strike is varying from western and northwestern to northly, with dip respectively to the south, southwest and west. The dip is gentle to medium.

### Mineralization in amphibole schist:

Pyrite and pyrrhotite may form a major part of the rock, especially in zones rich in quartz. Many of the outcrops are covered by rust. Thick layers of rust have sometimes made the examination of the rock difficult.

### Quartzite and granite

Quartzite and granite occur as lenses or layers in the amphibole schist. The layers can be more than 1 m in thickness. They are coarse, reddish and have no or few orientated minerals.

### Boulders

The boulders in the area are mostly of local rock types, and apparently of local origin. In the bog area to the northwest, there are several boulders of amphibolite, gneiss and granite. Some of these may be transported.

### Recommendations

Follow-up work in this area should consist of bedrock sampling of the sulphide zones.

AREA 46 Fig. 12

The bedrock in the area is well exposed.

The dominating rock type is amphibole schist which is bordering to granite in the northwest by fault zones.

Amphibole schist

The amphibole schist is foliated with mm-thick bands of green and light coloured minerals. It is mostly fine grained or occasionally medium grained. The main minerals are amphibole, quartz and feldspar, with various amounts of biotite. Varieties rich in biotite are mapped as mica schist. Much of the quartz in the amphibole schist occur as veins and lenses. Where these features are most prominent, the orientation of the rock is distorted. Outcrops mapped as quartzite are assumed to represent large quartz lenses in the amphibole schist. The direction of the strike is to the north-northeast with dip to the west-southwest. In the central part of the area, the strike is east-west dip to the south.

Mineralizations in amphibole schist:

Magnetite occur occasionally in some of the outcrops. Pyrite or pyrrhotite is found in various amounts in most of the outcrops. The highest amounts are found in outcrops that have a large content of quartz. In quartzite outcrops they can be a major constituent of the rock.

Most outcrops have rusty areas. In some cases the exposed bedrock is completely oxidised, and the minerals impossible to decide.

Granite


The granite is reddish, lacks orientation and is medium grained with quartz, feldspar and biotite as the main minerals.

Boulders

In the area there are boulders of amphibole schist, especially to the north. These are clearly of local origin.

Recommendations

Follow-up work in this area should consist of bedrock sampling of the sulphide zones.



LEGEND

Geology

-  Granite / P=pegmatite
-  Mica schist
-  Coarse, massive amphibolite or diabase
-  Tuff, tuffite
-  Medium to finegrained, often schistose and foliated, amphibolite
-  Carbonate rock, albite carbonate rock
-  Carbonate schist
-  Carbonate breccia
-  Sandstone, quartzite
-  Gneiss
-  Rust

-  Strike and dip (90 degrees division)
-  Rock boundary, certain and uncertain
-  Outcrop
-  Boulder
-  Group of boulders

- hem - hematite
  - py - pyrite
  - mt - magnetite
  - cp - chalcopyrite
- } As a major constituent (> 1/3) of the rock

Electromagnetics

-  Real comp. Horizontal loop 1777 Hz
-  Imag. comp. Horizontal loop 1777 Hz

Topography

-  Stream or river, lake
-  Bog
-  Elevation contour
-  Elevation 605
-  Path
-  Cart track
-  UTM grid with coordinates

AREA 45 and 46	M
	1:5000
Geological mapping EM survey	Målt: KB 83
	Tegn: KB 3/84
	Trace: HB 4/84
PROSPEKTERING A/S	Fig. 12



AREA 47 Fig. 13

The localization of the area is uncertain.

The greater part of the area is covered by bog. To the northeast and the southeast there are some outcrops. In the northeast there is an outcrop of foliated amphibolite and some outcrops of quartzite, rusty quartzite and one outcrop where nothing but rust can be seen. South of the quartzite there is an outcrop of granite, partly pegmatitic. There are layers of folded and foliated amphibole schist in the granitic rock mass. The strike of the rock is to the northeast and it has medium dip to the southeast.


In the southeast there is an area with folded layers of amphibolite schists and granitic rock. The rock also contains layers and masses of iron sulfides and is covered by rust in many places.

South of the amphibole-granite area, the rocks are gneissic.

Recommendations

Geological information on the area is sparse.

The anomaly is recommended investigated by till sampling.



LEGEND

Geology

- Granite / P=pegmatite
- Mica schist
- Coarse, massive, amphibolite or diabase
- Tuff, tuffite
- Medium to finegrained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust

- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3) of the rock

Electromagnetics

- Real comp. Horizontal loop 1777 Hz
- Imag. comp. Horizontal loop 1777 Hz

Topography

- Stream or river, lake
- Bog
- Elevation contour
- Elevation 605
- Path
- Cart track
- UTM grid with coordinates

Area 47 Lavvoai'vi Geological mapping EM survey	M
	1:5000
	Målt: KB 83
	Tegn: KB 3/84
Trace: HB 4/84	
PROSPEKTERING A/S	Fig.13

AREA 48, 49

AREA 48 Fig. 14

The area is completely covered by sand.

Recommendations

There is no information on the geology in this area.

Follow-up work should be deep till sampling.

AREA 49 Fig. 14


There are no outcrops in the area.

West of the area there is a boulder of amphibolite. East of the area there are a couple of boulders of mica schist.

Recommendations

There is little information on the geology in this area.

Recommended follow-up work is deep till sampling.



LEGEND

Geology

- Granite / P=pegmatite
- Mica schist
- Coarse, massive amphibolite or diabase
- Tuff, tuffite
- Medium to finegrained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust

- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcopyrite

} As a major constituent (> 1/3) of the rock

Electromagnetics

- Real comp. Horizontal loop 1777 Hz
- Imag comp. Horizontal loop 1777 Hz

Topography

- Stream or river, lake
- Bog
- Elevation contour
- Elevation 605
- Path
- Cart track
- UTM grid with coordinates

AREA 49 and 48 Lavvoai'vi Geological mapping EM survey	M
	1:5000
	Målt: KB 83
	Tegn: KB 3/84
Trace: HB 4/84	Fig. 14
PROSPEKTERING A/S	


AREA 50

AREA 50 Fig. 15

In this area there are a few outcrops of amphibolite.

Recommendations

Recommended follow-up work is deep till sampling.



LEGEND

Geology

- Granite / P=pegmatite
- Mica schist
- Coarse, massive amphibolite or diabase
- Tuff, tuffite
- Medium to finegrained, often schistose and foliated, amphibolite
- Carbonate rock, albite carbonate rock
- Carbonate schist
- Carbonate breccia
- Sandstone, quartzite
- Gneiss
- Rust
- Strike and dip (90 degrees division)
- Rock boundary, certain and uncertain
- Outcrop
- Boulder
- Group of boulders

hem - hematite  
 py - pyrite  
 mt - magnetite  
 cp - chalcocopyrite

} As a major constituent (> 1/3) of the rock

Electromagnetics

- Real comp. Horizontal loop 1777 Hz
- Imag. comp. Horizontal loop 1777 Hz

Topography

- Stream or river, lake
- Bog
- Elevation contour
- Elevation 605
- Path
- Cart track
- UTM grid with coordinates

AREA 50 Lavvoai'vi Geological mapping EM survey	M
	1:5000
	Målt: KB 83
	Tegn: KB 3/84
	Trace: HB 4/84
PROSPEKTERING A/S	Fig. 15